To 2020 NASA TM-75598

M-75598 ARchwod

THE EFFECTS OF STRONG SHOCK WAVES ON MORTALITY RATES AND PERCENTAGES OF PULMONARY LESIONS IN RATS AS A FUNCTION OF THE NUMBER OF EXPOSURES

Patrick Vassout and Georges Parmentier and Planeer (in the buins)

der for in the same

17 4 mini To 2 min

Translation of "Etude des effets des ondes de choc fortes sur les taux de letalite et les pourcentages de lesions pulmonaires chez le rat en fonction du nombre d'expositions", Institut franco-allemend de recherches (I.S.L.) Sant-Louis (France), Unpublished report, 1978, pp. 1-13.

STRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

20000912 103

Our best for 100 grat from Bowens report in LDso 48,6 ± 1.3 pm.

Our Hereshollfor LH C.67 born

ISL 11 11 LIF 0.5 born

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION WASHINGTON, D.C. 20546 DECEMBER 1978

Reproduced From Best Available Copy

Die Ought and State

THE EFFECTS OF STRONG SHOCK WAVES ON MORTALITY RATES AND PERCENTAGES
OF PULMONARY LESIONS IN RATS AS A FUNCTION OF THE NUMBER OF
EXPOSURES

Patrick Vassout and Georges Parmentier

Introduction

The effects of strong blast waves have been studied in particular */
since the Second World War. In general, authors have described the
relationships which exist between such and such a parameter of the
blast wave and the percentages of pulmonary or intestinal lesions,
as well as the mortality rates, observed in subjects of a given
species for a single exposure.

The literature reviews in [1] and [2] show that so far the parameter "number of exposures" has not been dealt with by the several investigators in question. Indeed, it would not seem very realistic to imagine that an individual could be subjected to a succession of pure blasts. The appearance of new weapons based on the principle of gas detonation, as well as the development of weapons which produce very high overpressure levels at the start of the blast pose the problem concerning the effect of the number of exposures to a strong blast wave on the extent of pulmonary lesions and on mortality rates.

We have therefore undertaken to study the "number of exposures" parameter on pulmonary lesions and mortality rates in the rat [3].

1. Experimental Apparatus

The experimental animals were female rats with an average body weight of 100 g. Each animal was placed in a wire mesh cage [1]. The cages were placed symmetrically around the explosive charge. With this arrangement it was possible to apply the same blast wave to each animal while changing as little as possible the pressure profile of the wave.

The explosive used was "plastit" made by Dottikom Co., Switzerland.

^{*}Numbers in the margin indicate pagination in the foreign text.

1.1. Determination of the Amounts of Explosive Required

By applying the laws of similarity valid for spherical explosives [4] [5], we determined the amounts of explosive necessary to obtain the desired overpressures while maintaining a constant duration of the positive first phase t_0 .

1.2. Recording Pressure Changes

The pressure signals (Fig. 2) were recorded by means of KISTLER / 6031 piczoelectric dynamic pick-ups placed in the tapered waves. These pick-ups were connected to BIOMATION 805 transistorized analog digital signal recorders. The pick-ups were installed at the same distance from the explosive charge as the experimental animals and measured the pressure in free space. The animals, placed on [word illegible], were exposed to the incident pressure and the pressure reflected by the ground. The pressure levels discussed here always refer to those which existed at the level of the animals. (Over-pressure measured in free space multiplied by the reflection factor)

2. Results

2.1. Mortality Rates as a Function of the Number of Exposures 2.1.1. Determination of the Median Lethal Dose for Rats

The median lethal dose (LD $_{50}$) is the crest overpressure which for a given length of time T $_{0}$ for the incident blast wave causes death in 50% of the animals exposed at the end of a given length of time after exposure, for example 24 hours.

Based on the studies of Richmond and White (1962) [6], done on many species of animals, we can estimate the crest overpressure corresponding to the LD_{50} for rats for a given to of the blast wave (LD_{50} = 3 bars for to = 2 ms). We have chosen as [word illegible] of the blast wave the value to = 2 ms. Indeed, according to these authors, beyond this length of time this parameter no longer has any effect on the mortality rate in rats.

King to the state of the

For this 2-ms length of time we have studied the mortality rate in rats for a single exposure to an overpressure blast wave between

3 and 5 bars (Fig. 3). It should be noted that all of the overpressure less than 3 bars did not cause any mortality in our upper experiments.

It appears that the LD₅₀ for rats, 24 hours after a single (0 exposure, is located around 3.5 bars. We also find that the difference in overpressure necessary to go from a mortality rate of 10% to one of 90% five minutes after exposure is 1.5 bars in rats. $\chi_{(4,3,4)} = 2.36 f^{-2}$

2.1.2. Effect of the Number of Exposures on Rat Mortality Rates 74

For the purpose of determining relationships between mortality rates, the number of exposures and crest overpressure, series of 40 animals were exposed to blast waves of t = 2 s produced at regular 4-minute intervals.

Fig. 4 shows the results obtained in the overpressure range between 2 and 4 bars. The dotted line curve which represents the LD₅₀ reveals that a twofold increase in the level of the crest overpressure is offset by a 20-fold decrease in the number of exposures.

2.2. Percentages of Pulmonary Lesions Due to Exposure to Strong Blast Waves

2.2.1. Definition of Lesion Criteria Used in our Experiments

Fig. 3 shows that a single exposure to a wave overpressure blast wave less than 3 bars does not cause any mortality. However, autopsy reveals that the animals have hemorrhagic pulmonary lesions ranging from simple petechia to a completely hemorrhagic lobe. We have taken as the lesion threshold the small hemorrhage isolated on a single lobe according to the criteria proposed by Richmond et al. [7].

2.2.2. Determination of the Overpressure which Causes Pulmonary Lesions in 50% of the Experimental Animals

As in previous experiments, we first of all studied the influence of a single exposure to varied crest overpressures on the rate of pulmonary lesions.

The overpressure which causes the appearance of pulmonary lesions in 50% of the experimental animals is situated around 500 mbars (Fig. 5). An overpressure of 2.5 bars lasting for the same amount of time (2 ms) causes lesions to appear in all of the animals exposed.

/5

2.2.3. Effect of the Number of Exposures on the Percentage of Animals Showing Pulmonary Lesions

Groups of 40 animals were exposed to series of blast waves all lasting t_0 = 2 ms and of overpressures ranging between 32 and 500 mbars. The results obtained are plotted in Fig. 6. Note that for a constant percentage of animals with lesions (dotted line curve), a two-fold increase in the number of exposures must be compensated for by a quartering of the crest overpressure. This result shows a different relationship between the wave overpressure and the number of exposures experienced from [word illegible] observed on the basis of mortality rates.

3. Trial Extrapolation for a Theoretical Animal Weighing 70 kg and Conclusion

Many authors such as Zuckerman 1941 [10], White 1959 [9] Richmond 1968 [7] and Jones 1969 [9] think that it is possible to extrapolate the results obtained in animals to man. They indicate that the LD $_{50}$ applicable to man would be on the order of 10 bars for a blast wave lasting 2 ms (t_{0}). These authors also think that the threshold for the appearance of pulmonary lesions in man surround one-fourth of the LD $_{50}$, or around 2.5 bars.

As for the effect of the number of exposures on mortality rate, extrapolations of our results to man indicates that twenty exposures to a crest overpressure of 5 bars lasting 2 ms (t_0) experienced at intervals on the order of 5 [word illegible] would cause death in 50% of the subjects exposed.

As for the effect of the number of exposures on the appearance of pulmonary lesions, we fear that lesions might appear in man for crest overpressure values plainly less than those currently accepted (2.5 bars for a too f [word missing] where the subject is exposed to a series of strong blast waves).

+ Fletcher # from Bowens report

11




Fig. 1. Arrangement of Rat Cages





Fig. 2. Typical recording of pressure signals for an explosive mass.

and the second sections of the second constitution of the second	une exposition	n 24 h d'ani.	3.0	-	83 68	49 23	5	91 35	
OF BITTE CHEST SECURIOR PETERS DEBAS	après ur	2 h 4 h	S. S	- -	38			83 · 88	
A laing and an amendadem taken taken and an ann an	ips écoulé	39mm 4 h	(a)	+	2	ເລ	25	3.	9
AN	Temps	[bar] Smn	0	Ef ev	_	100 100 100	S. S. S.	_ _	ි වා
Distance		[cm]	278	270	105.17	8.76.67	267	6	N 50 N
Poids	۸ مان مان مان مان مان	[5]	2,2 16 13.88	0000	2000	13/	3,416 1560	best	

Fig. 3. Mortality rates observed after an exposure to blast waves of variable Δe and constant t_{o} (2 ms) in rats weighing 100 g.

Key:

Weight of "plastit" in grams Elapsed time after explosion Number of animals မှ ကုပ်



Effect of the number of blast waves experienced as a function of overpressure with a constant t_0 (2 ms) on mortality rate in rats.

Number of blast waves Key: a.

b.


 $^{\mathrm{LD}_{50}}_{\mathrm{Mortality}}$ rate c.

24 hours after exposure

Par De l'annie de l'an


Fig. 5. [Caption illegible]

Key; a. Percent of animals with lesions



Rigaud, P., "Effect of Blast Waves on Living Creatures in

- 1. General." Paper presented to the Group of French-Speaking Acousticians (GALF) in May 1970. Notice ISL NB 2/71, [1971).
- 2. Rigaud, P. and A. Dancer, "Effect of Blast Waves on Living Creatures," Notice ISL NB 6/73, (1973).
- Vassout, P., G. Paramentier and A. Dancer, "The Effect of 3. Strong Blast Waves on the Mortality Rate and the Appearance of Pulmonary Lesions in Rats," ISL Report R 922/78, (1978).
- 4. Froböse, M., "Propagation Functions of Spherical Aerial Blast Waves," ISL Report R 3/68 (1968).
- Froböse, M. and O. Wünsche, "Mortality of Albino Rats Exposed 5. to an Aerial Blast Wave as a Function of the Direction of the Blast and the Shape of the Pressure Curve," ISL Report R 2/59, (1959).
- 6. [No No. 6 listed]
- Richmond, D.R., E.G. Damon, G.I. Bowen, E. Fletcher and C.S. White, 7. "Air Blast Sudies with Eight Species of Mammals," Lovelace Foundation for Medicine Education and Research, New York, USA 1 (1966) DASA 1854. "The Relationship Between Selected Blast Wave Parameters on the Response of Mammals Exposed to Air Blast," A. of New York 1 Acad. of Sciences 152, 103-121, (1968). "A Tentative Estimation of Man's Tolerance to Overpressures From Air Blast," Lovelace Foundation for Medicine Education and Research, New York, USA (1962) DASA 1335.
- White, C.S. and D. R. Richmond, "Blast Biology," Lovelace 8. Foundation for Medical Education and Research, New Mexico, T.I.D. 5764, (1959).
- 9. Jones, R.D., D.R. Richmond, and R.E. Flechter, "A Reappraisal of Man's Tolerance to Indirect Air Blast Injury," Lovelace Foundation for Medicine Education and Research, New Mexico, USA, (1969).
- Zuckerman, S., "Experimental Study of Blast Injury to the Lungs," 10. Lancet II, pp. 219-238, (1940).




Effect of the Number of Blast Waves Experienced 0.9mi 0.469 as a Function of the Overpressure of 2-Msec Duration on the Percentage of Rats with Lesions. Figure 4.

154

pai in 1/50 1/ the 6050 god.

sheep 1950 01 1 tens, the 20 was 160 ... say 65 than 1/50 of the 113 pai our would expect here is if Topsi de 15 pm



Effect of the Number of Blast Waves Experienced as a Function of Overpressure with a Constant 2-Msec Duration on Mortality Rate in Rats. Figure 3.

154

1. Report No. NASA TM-75598	2. Government Accession No.		3. Recipient's Catalog No.						
4. Title and Subtitle THE EFFECTS OF STRONG TALITY RATES AND PERC LESIONS IN RATS AS A 7. Author(s) OF EXPOSURES Patrick Vassout and Institut of Franco Go	6. Performing Organization Code ER 8. Performing Organization Report No.								
Sant-Louis, France 9. Performing Organization Name and Leo Kanner Associates Redwood City, Califor		11. Contract or Grant No. NASW-3199 13. Type of Report and Period Covered							
12. Sponsoring Agency Name and Addre National Aeronautics tration, Washington,	Translation 14. Sponsoring Agency Code								
Translation of "Etude des effets des ondes de choc fortes sur les taux de letalite et les pourcentages de lesions pulmonaires chez le rat en fonction du nombre d'expositions", Institut franco-allemend de recherches (I.S.L.) Sant-Louis (France), Unpublished report, 1978, pp. 1-13.									
This study of pulmonary lesions and mortality rates caused in rats by exposure to strong shock waves of constant duration revealed that with regard to the pulmonary lesions, twice the number of exposures is compensated for by quartering the overpressure of the wave crest, and with regard to the mortality rates, it revealed that halving the overpressure of the wave crest is offset by a 20-fold increase in the number of exposures.									
17. Key Words (Selected by Author(s))		Unclassified Unlimited							
19. Security Classif. (of this report) Unclassified	20. Security Class Unclass		21. No. of Pages	22. Price					